



Regular Articles

Dynamics of composition operators on function spaces defined by local and global properties

Angela A. Albanese^a, Enrique Jordá^{b,*}, Claudio Mele^a^a Dipartimento di Matematica e Fisica "E. De Giorgi", Università del Salento, C.P. 193, I-73100 Lecce, Italy^b Instituto Universitario de Matemática Pura y Aplicada IUMPA, Universitat Politècnica de València, EPS Alcoy Plaza Ferrándiz y Carbonell, s/n, 03801 Alcoy (Alicante), Spain

ARTICLE INFO

Article history:

Received 15 December 2021
Available online 4 May 2022
Submitted by J. Bonet

Keywords:

Spaces of continuous functions
Composition operator
Power bounded operator
Mean ergodic operator
Hypercyclic operator
Supercyclic operator

ABSTRACT

In this paper we consider composition operators on locally convex spaces of functions defined on \mathbb{R} . We prove results concerning supercyclicity, power boundedness, mean ergodicity and convergence of the iterates in the strong operator topology.

© 2022 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

In this paper we deal with composition operators defined in locally convex Hausdorff spaces of real valued continuous functions. We study when the operator has *big orbits*, i.e., when it is supercyclic or hypercyclic, and also when it has *small orbits*, i.e., when the operator is power bounded or mean ergodic. We collect in this section our results and relate them with recent literature.

We focus our attention on composition operators on locally convex Hausdorff spaces $X \hookrightarrow C^m(\mathbb{R})$, for $m \in \mathbb{N}_0 \cup \{\infty\}$. For a function $\varphi \in C^m(\mathbb{R})$, being increasing and with no fixed points is equivalent to being *strongly runaway*, i.e., for all compact subsets $K \subset \mathbb{R}$ there is $n_0 \in \mathbb{N}$ such that $\varphi^n(K) \cap K = \emptyset$ for all $n \geq n_0$ (see [20, Lemma 4.1] for a proof). Kalmes got in [18, Corollary 4.1, Corollary 4.2] the equivalence between hypercyclicity and strongly runaway with no vanishing derivative in a more general context, which includes weighted composition operators defined in function spaces of several variables. The results of Kalmes generalize those of Przewacki obtained in [20, Lemma 4.1, Theorem 4.5, Theorem 4.6]

* Corresponding author.

E-mail addresses: angela.albanese@unisalento.it (A.A. Albanese), ejorda@mat.upv.es (E. Jordá), claudio.mele1@unisalento.it (C. Mele).

for composition operators on $C^\infty(\Omega)$, with $\Omega \subseteq \mathbb{R}^d$ open. Our main result here is that, under natural assumptions on X , C_φ being weakly supercyclic implies that φ is increasing, with no vanishing derivative when $m > 0$, and without fixed points. This, using previous works of Bonet and Domański and Kalmes, permits us to conclude that weak supercyclicity is equivalent to being mixing for composition operators defined on $C^m(\mathbb{R})$ with $m \in \mathbb{N} \cup \{\infty\}$ and that weak supercyclicity implies to be mixing for composition operators defined on $\mathcal{A}(\mathbb{R})$.

We also obtain results for power boundedness and mean ergodicity of composition operators. We prove that for a Montel space $X \hookrightarrow C^1(\mathbb{R})$ satisfying that, for every symbol $\varphi \in X$

(*) The composition operator $C_\varphi : X \rightarrow X$ is power bounded if, and only if, the sequence $\{\varphi_n\}_{n \in \mathbb{N}}$ of iterates of φ is bounded in X ,

under natural assumptions on X , choosing a monotonic $\varphi \in X$ with $\varphi_2(x) \neq x$, then the composition operator C_φ is power bounded if, and only if, φ has a unique fixed point a to which the sequence of the iterates φ_n of φ converges if, and only if, the sequence $\{C_{\varphi_n}\}_{n \in \mathbb{N}}$ is convergent in $\mathcal{L}_s(X)$ to C_a ($C_a : X \rightarrow X, f \mapsto f(a)$).

Furthermore, if we restrict to particular locally convex spaces X and a polynomial $\varphi(x)$, then the composition operator is equivalently power bounded and mean ergodic if, and only if, $\varphi(x) = ax + b$, $|a| < 1$. The excluded cases $\varphi(x) = x$ and $\varphi(x) = -x + b$ produce trivially power bounded uniformly mean ergodic composition operators. The equivalence between power boundedness, mean ergodicity and the convergence of the iterates of φ to the unique fixed point of φ has been proved in [9] to be true for $\mathcal{A}(\mathbb{R})$, with no restriction on the symbol. Besides $\mathcal{A}(\mathbb{R})$, our result applies to $C^\infty(\mathbb{R})$, which satisfies (*) by [17] and to the space $\mathcal{O}_M(\mathbb{R}) := \text{proj}_{\overline{m}} \text{ind}_{\overline{n}} \mathcal{O}_n^m(\mathbb{R})$ of the multipliers of the classical space $\mathcal{S}(\mathbb{R})$.

The structure of the paper is the following: Section 2 is devoted to preliminaries and to the study of the symbols in $\mathcal{O}_M(\mathbb{R})$ and $\mathcal{O}^m(\mathbb{R})$, $m \in \mathbb{N}$. In Section 3 we study weakly supercyclic composition operators. In Section 4 we obtain respectively the results on dynamics explained above. Finally, Section 5 is devoted to show that $\mathcal{O}_M(\mathbb{R})$ and $\mathcal{O}^m(\mathbb{R})$, $m \in \mathbb{N}$, satisfy condition (*).

2. Preliminaries and the action of the composition operator on $\mathcal{O}_M(\mathbb{R})$

2.1. Preliminaries

Through the paper, we denote by \mathbb{N} the set of all positive integer numbers and set $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$.

Let X be a locally convex Hausdorff space (briefly, lchS). We denote by $\mathcal{L}(X)$ the space of all continuous linear operators from X into itself.

An operator $T \in \mathcal{L}(X)$, with X a lchS, is called *power bounded* if $\{T^n\}_{n \in \mathbb{N}}$ is an equicontinuous subset of $\mathcal{L}(X)$.

The Cesàro means of an operator $T \in \mathcal{L}(X)$, with X a lchS, are defined by

$$T_{[n]} := \frac{1}{n} \sum_{m=1}^n T^m, \quad n \in \mathbb{N}.$$

The operator T is called *mean ergodic* (resp. *uniformly mean ergodic*) if $\{T_{[n]}\}_{n \in \mathbb{N}}$ is a convergent sequence in $\mathcal{L}_s(X)$ (resp. in $\mathcal{L}_b(X)$), where $\mathcal{L}_s(X)$ denotes $\mathcal{L}(X)$ endowed with the strong operator topology τ_s (resp. $\mathcal{L}_b(X)$ denotes $\mathcal{L}(X)$ endowed with the topology τ_b of the uniform convergence on bounded subsets of X).

For reflexive Fréchet spaces (in Montel Fréchet spaces, resp.) every power bounded operator is necessarily mean ergodic (uniformly mean ergodic, resp.) by [1, Corollary 2.7, Proposition 2.9]. There exist mean ergodic operators which are not power bounded, see, f.i., [16, §6]. For further results on mean ergodic operators we

refer to [19,22]. For recent results on mean ergodic operators in lcHs' we refer to [1–4,7,12,17], for example, and the references therein.

Given an operator $T \in \mathcal{L}(X)$, we denote by $\sigma(T)$ the *spectrum* of T , i.e., the subset of \mathbb{C} formed by the λ such that $\lambda I - T$ is not invertible, and by $\sigma_p(T)$ its *point spectrum*, i.e., the subset of $\sigma(T)$ formed by those λ such that $\lambda I - T$ is not injective.

Moreover, we recall that T' denotes as usual the dual operator of T . We have that $T \in \mathcal{L}(X'_\beta)$ and also $T' \in \mathcal{L}(X'_\sigma)$, where X'_β and X'_σ stand for the topological dual of X endowed with the strong topology $\beta(X', X)$ and the weak topology $\sigma(X', X)$ resp.

Let X be a lcHs of functions defined on \mathbb{R} and let $\varphi: \mathbb{R} \rightarrow \mathbb{R}$ be a function. If $f \circ \varphi \in X$ for all $f \in X$, then we can consider the composition operator $C_\varphi: X \rightarrow X, f \mapsto f \circ \varphi$. The operator C_φ is clearly linear. In case that $C_\varphi \in \mathcal{L}(X)$, the function φ is said to be a *symbol* for X .

Given a function $\varphi: \mathbb{R} \rightarrow \mathbb{R}$, we denote by φ_n the n -th iterate of φ for all $n \in \mathbb{N}$, i.e., $\varphi_n := \varphi \circ \dots \circ \varphi$ n -times, and define $\varphi_{[n]} := \frac{1}{n} \sum_{k=1}^n \varphi_k$ for all $n \in \mathbb{N}$. In case the function φ is bijective and hence there exists $\varphi^{-1}: \mathbb{R} \rightarrow \mathbb{R}$, we denote by φ_{-n} the n -th iterate of φ^{-1} for all $n \in \mathbb{N}$. We also recall that a function $\varphi: \mathbb{R} \rightarrow \mathbb{R}$ is said to be increasing (decreasing, resp.) if $\varphi(x) < \varphi(y)$ ($\varphi(x) > \varphi(y)$, resp.) for every $x, y \in \mathbb{R}$ such that $x < y$. The function φ is said to be non decreasing (non increasing, resp.) if $\varphi(x) \leq \varphi(y)$ ($\varphi(x) \geq \varphi(y)$, resp.) for every $x, y \in \mathbb{R}$ such that $x < y$.

In what follows, we recall the necessary definitions and some basic properties of the spaces $\mathcal{S}(\mathbb{R})$ and $\mathcal{O}_M(\mathbb{R})$. For general information about function spaces in the theory of distributions we refer to [11,21].

The space $\mathcal{S}(\mathbb{R})$ of *rapidly decreasing functions* is a nuclear Fréchet space and hence, it is Montel and reflexive. Accordingly, its strong dual $\mathcal{S}'(\mathbb{R})$ is a nuclear lcHs. In particular, $\mathcal{S}'(\mathbb{R})$ is a barreled and bornological lcHs.

The space $\mathcal{O}_M(\mathbb{R})$ of *slowly increasing functions* on \mathbb{R} is given by

$$\mathcal{O}_M(\mathbb{R}) = \bigcap_{m=1}^\infty \bigcup_{n=1}^\infty \mathcal{O}_n^m(\mathbb{R}),$$

where $\mathcal{O}_n^m(\mathbb{R}) := \{f \in C^m(\mathbb{R}) : |f|_{m,n} := \sup_{x \in \mathbb{R}} \sup_{0 \leq i \leq m} (1+x^2)^{-n} |f^{(i)}(x)| < \infty\}$, endowed with the norm $|\cdot|_{m,n}$, is a Banach space for any $m, n \in \mathbb{N}$. The space $\mathcal{O}_M(\mathbb{R})$, endowed with its natural lc-topology, i.e., $\mathcal{O}_M(\mathbb{R}) := \text{proj}_{\leftarrow} \text{ind}_{\rightarrow} \mathcal{O}_n^m(\mathbb{R})$, is a projective limit of the complete (LB)-spaces $\mathcal{O}^m(\mathbb{R}) := \text{ind}_{\rightarrow} \mathcal{O}_n^m(\mathbb{R})$, for $m \in \mathbb{N}$. In particular, $\mathcal{O}_M(\mathbb{R})$ is a bornological nuclear lcHs (hence, Montel and reflexive), and it is continuously embedded in $C^\infty(\mathbb{R})$ (see [15]). Furthermore, a fundamental system of continuous norms on $\mathcal{O}_M(\mathbb{R})$ is given by

$$p_{m,v}(f) = \sup_{x \in \mathbb{R}} \sup_{0 \leq i \leq m} |v(x)| |f^{(i)}(x)|, \quad f \in \mathcal{O}_M(\mathbb{R}),$$

where $v \in \mathcal{S}(\mathbb{R})$ and $m \in \mathbb{N}$ (see, f.i., [11]).

The space $\mathcal{O}_M(\mathbb{R})$ is the space of multipliers of $\mathcal{S}(\mathbb{R})$ and its strong dual $\mathcal{S}'(\mathbb{R})$. So, for any fixed $h \in \mathcal{O}_M(\mathbb{R})$, the multiplication operator $M_h: \mathcal{S}(\mathbb{R}) \rightarrow \mathcal{S}(\mathbb{R}), f \mapsto hf$, is continuous.

The spaces $C^m(\mathbb{R}), m \in \mathbb{N}_0 \cup \{\infty\}$, are always endowed with their natural lc-topology τ_c given from the canonical fundamental sequence of seminorms. We point out that $(C^m(\mathbb{R}), \tau_c)$ is a Fréchet space. In particular, $(C^\infty(\mathbb{R}), \tau_c)$ is a nuclear Fréchet space. Moreover, we denote by $\mathcal{D}^m(\mathbb{R}) := \{f \in C^m(\mathbb{R}) : \text{supp} f \text{ is compact}\}$, for $m \in \mathbb{N}_0 \cup \{\infty\}$ (we write $\mathcal{D}(\mathbb{R})$ for $m = \infty$). Then $\mathcal{D}^m(\mathbb{R}) = \bigcup_{k=1}^\infty C_0^m([-k, k])$ for all $m \in \mathbb{N}$ and so, endowed with its natural lc-topology, i.e., $\mathcal{D}^m(\mathbb{R}) := \text{ind}_{\leftarrow} C_0^m([-k, k])$, is a complete (LB)-space ((LF)-space, for $m = \infty$).

We remark the following fact.

Proposition 2.1. *Let $m \in \mathbb{N}$ and let B be a bounded subset of $\mathcal{O}^m(\mathbb{R})$. Then the spaces $\mathcal{O}^m(\mathbb{R})$ and $C^m(\mathbb{R})$ induce on B the same topology.*

Proof. For any $p \in \mathbb{N}$ we will show that the spaces $\mathcal{O}_{p+1}^m(\mathbb{R})$ and $C^m(\mathbb{R})$ induce the same topology on the unit ball B_p^m of $\mathcal{O}_p^m(\mathbb{R})$. To do this, we only need to show that for each $\varepsilon > 0$, $U := \{f \in B_p^m : |f|_{m,p+1} < \varepsilon\}$ is a neighborhood of 0 in B_p^m endowed with the topology inherited from $C^m(\mathbb{R})$. So, let $M > 0$ be such that $(1+x^2)^{-1} < \varepsilon$, whenever $|x| > M$ and let $V := \{f \in C^m(\mathbb{R}) : |f^{(i)}(x)| < \varepsilon \text{ for all } i = 0, \dots, m, |x| \leq M\}$. Then $B_p^m \cap V \subseteq U$. Indeed, if $f \in B_p^m \cap V$, then

$$(1+x^2)^{-p-1}|f^{(i)}(x)| \leq |f^{(i)}(x)| < \varepsilon, \quad i = 0, \dots, m, |x| \leq M,$$

and

$$(1+x^2)^{-p-1}|f^{(i)}(x)| = (1+x^2)^{-p}|f^{(i)}(x)|(1+x^2)^{-1} \leq (1+x^2)^{-1} < \varepsilon, \quad i = 0, \dots, m, |x| > M.$$

Therefore, $f \in U$. The conclusion follows from the fact that the bounded subsets of $\mathcal{O}^m(\mathbb{R})$ are localized, i.e., $\mathcal{O}^m(\mathbb{R})$ is a regular (LB)-space. \square

Remark 2.2. As a consequence of Proposition 2.1, we get that a sequence $\{\varphi_n\}_{n \in \mathbb{N}} \subset \mathcal{O}^m(\mathbb{R})$ is convergent in $\mathcal{O}^m(\mathbb{R})$ to some $\varphi \in \mathcal{O}^m(\mathbb{R})$ if, and only if, $\{\varphi_n\}_{n \in \mathbb{N}}$ is a bounded sequence in $\mathcal{O}^m(\mathbb{R})$ and $\lim_{n \rightarrow \infty} \varphi_n = \varphi$ in $C^m(\mathbb{R})$. Therefore, as $\mathcal{O}_M(\mathbb{R})$ is the projective limit of the complete (LB)-spaces $\mathcal{O}^m(\mathbb{R})$, for $m \in \mathbb{N}$, we can then conclude that a sequence $\{\varphi_n\}_{n \in \mathbb{N}} \subset \mathcal{O}_M(\mathbb{R})$ is convergent in $\mathcal{O}_M(\mathbb{R})$ to some $\varphi \in \mathcal{O}_M(\mathbb{R})$ if, and only if, $\{\varphi_n\}_{n \in \mathbb{N}}$ is a bounded sequence in $\mathcal{O}_M(\mathbb{R})$ and $\lim_{n \rightarrow \infty} \varphi_n = \varphi$ in $C^\infty(\mathbb{R})$. Therefore, for every sequence $\{f_n\}_{n \in \mathbb{N}} \subset \mathcal{D}(\mathbb{R})$ satisfying the condition for every compact subset K of \mathbb{R} there exists $n_0 \in \mathbb{N}$ such that $\text{supp} f_n \cap K = \emptyset$ for all $n \geq n_0$, we get that $\{f_n\}_{n \in \mathbb{N}}$ is convergent to 0 in $\mathcal{O}^m(\mathbb{R})$ (in $\mathcal{O}_M(\mathbb{R})$, resp.) if, and only if, it is a bounded sequence of $\mathcal{O}^m(\mathbb{R})$ (of $\mathcal{O}_M(\mathbb{R})$, resp.).

We point out that there are examples of sequences $\{f_n\}_{n \in \mathbb{N}} \subset \mathcal{D}(\mathbb{R})$ satisfying the previous condition but not bounded in $\mathcal{O}^m(\mathbb{R})$ (in $\mathcal{O}_M(\mathbb{R})$, resp.). Indeed, for all $n \in \mathbb{N}$ let $f_n \in \mathcal{D}(\mathbb{R})$ be such that $f_n(x) := n(1+x^2)^n$ for $x \in K_n := [n + \frac{1}{n}, n + 1 - \frac{1}{n}]$ and $\text{supp} f_n \subseteq [n, n + 1]$. Then for any $m, p \in \mathbb{N}$ we have

$$\begin{aligned} |f_n|_{m,p} &= \sup_{x \in \mathbb{R}} \sup_{0 \leq i \leq m} (1+x^2)^{-p} |f_n^{(i)}(x)| \\ &\geq \sup_{x \in K_n} (1+x^2)^{-p} |f_n(x)| = n \sup_{x \in K_n} \frac{(1+x^2)^n}{(1+x^2)^p} \geq n, \quad \text{for all } n \geq p. \end{aligned}$$

This means that the sequence $\{f_n\}_{n \in \mathbb{N}}$ cannot be bounded in the Banach space $\mathcal{O}_p^m(\mathbb{R})$ for all $p \in \mathbb{N}$ and hence in $\mathcal{O}^m(\mathbb{R})$, being $\mathcal{O}^m(\mathbb{R}) = \text{ind}_{\mathbb{R}} \mathcal{O}_p^m(\mathbb{R})$ a regular (LB)-space.

2.2. The composition operator on $\mathcal{O}_M(\mathbb{R})$

In [13] the symbols $\varphi \in C^\infty(\mathbb{R})$ for $\mathcal{S}(\mathbb{R})$ have been characterized. Precisely, [13, Theorem 2.3] states that a function $\varphi \in C^\infty(\mathbb{R})$ is a symbol for $\mathcal{S}(\mathbb{R})$ if, and only if, the following conditions are satisfied:

(1) For all $j \in \mathbb{N}_0$ there exist $C, p > 0$ such that, for every $x \in \mathbb{R}$

$$|\varphi^{(j)}(x)| \leq C(1 + \varphi(x)^2)^p;$$

(2) There exists $k > 0$ such that

$$|\varphi(x)| \geq |x|^{\frac{1}{k}}, \quad \forall |x| \geq k.$$

We now characterize the function $\varphi \in C^\infty(\mathbb{R})$ such that C_φ acts continuously from $\mathcal{O}_M(\mathbb{R})$ into itself. In order to do this, we first establish the following result.

Theorem 2.3. *Let $m \in \mathbb{N}$ and let $\varphi \in C^m(\mathbb{R})$. Then the following properties are equivalent:*

- (1) C_φ acts continuously from $\mathcal{O}^m(\mathbb{R})$ into itself;
- (2) $\varphi \in \mathcal{O}^m(\mathbb{R})$.

Proof. (1) \Rightarrow (2). Consider the function $p_1(x) := x$, for $x \in \mathbb{R}$, which belongs to $\mathcal{O}^m(\mathbb{R})$. Then $\varphi = C_\varphi p_1 \in \mathcal{O}^m(\mathbb{R})$. So, (2) is satisfied.

(2) \Rightarrow (1). We claim that for all $n \in \mathbb{N}$ there exists $n' \in \mathbb{N}$ such that $C_\varphi : \mathcal{O}_n^m(\mathbb{R}) \rightarrow \mathcal{O}_{n'}^m(\mathbb{R})$ is continuous. Indeed, for fixed $n \in \mathbb{N}$ and $f \in \mathcal{O}_n^m(\mathbb{R})$, we observe for every $x \in \mathbb{R}$ and $i = 0, \dots, m$ that

$$(C_\varphi f)^{(i)}(x) = \sum \frac{i!}{k_1!k_2! \dots k_n!} f^{(k)}(\varphi(x)) \left(\frac{\varphi'(x)}{1!}\right)^{k_1} \left(\frac{\varphi''(x)}{2!}\right)^{k_2} \dots \left(\frac{\varphi^{(i)}(x)}{i!}\right)^{k_i}, \tag{2.1}$$

where the sum is extended over all $(k_1, k_2, \dots, k_i) \in \mathbb{N}_0^i$ such that $k_1 + 2k_2 + \dots + ik_i = i$ and $k_1 + k_2 + \dots + k_i = k$ (observe that $k \leq i$). Since by assumption on φ , i.e., $\varphi \in \mathcal{O}^m(\mathbb{R})$, there exist $D > 0$ and $p \in \mathbb{N}$ such that $|\varphi^{(j)}(x)| \leq D(1 + x^2)^p$ for $x \in \mathbb{R}$ and $j = 0, \dots, m$ and hence $(1 + \varphi(x)^2) \leq D'(1 + x^2)^{2p}$ for $x \in \mathbb{R}$ and a suitable constant $D' > 0$, from (2.1) it follows for every $x \in \mathbb{R}$ and $i = 0, \dots, m$ that

$$\begin{aligned} |(C_\varphi f)^{(i)}(x)| &\leq \sum \frac{i!}{k_1!k_2! \dots k_i!} |f^{(k)}(\varphi(x))| \left|\frac{\varphi'(x)}{1!}\right|^{k_1} \left|\frac{\varphi''(x)}{2!}\right|^{k_2} \dots \left|\frac{\varphi^{(i)}(x)}{i!}\right|^{k_i} \\ &\leq \sum \frac{i!}{k_1!k_2! \dots k_i!} |f|_{m,n} (1 + \varphi(x)^2)^n \frac{D^k (1 + x^2)^{pk}}{1!^{k_1} 2!^{k_2} \dots i!^{k_i}} \\ &\leq C_i |f|_{m,n} (1 + x^2)^{p(2n+i)} \leq C |f|_{m,n} (1 + x^2)^{p(2n+m)}, \end{aligned}$$

where $C_i := (D')^n \sum \frac{i!}{k_1!k_2! \dots k_i!} \frac{D^k}{1!^{k_1} 2!^{k_2} \dots i!^{k_i}}$ and $C := \max_{i=0, \dots, m} C_i$. Accordingly, if $n' := p(2n + m) \in \mathbb{N}$, we get that

$$|C_\varphi f|_{m,n'} \leq C |f|_{m,n}.$$

Since $n \in \mathbb{N}$ and $f \in \mathcal{O}_n^m(\mathbb{R})$ are arbitrary, the claim is proved. \square

As a consequence of the result above, we obtain the class of symbols for which $C_\varphi \in \mathcal{L}(\mathcal{O}_M(\mathbb{R}))$.

Theorem 2.4. *Let $\varphi \in C^\infty(\mathbb{R})$. Then the following properties are equivalent:*

- (1) C_φ acts continuously from $\mathcal{O}_M(\mathbb{R})$ into itself;
- (2) C_φ acts continuously from $\mathcal{O}^m(\mathbb{R})$ into itself for all $m \in \mathbb{N}$;
- (3) $\varphi \in \mathcal{O}_M(\mathbb{R})$.

Proof. (1) \Rightarrow (3). Consider the function $p_1(x) := x$, for $x \in \mathbb{R}$, which belongs to $\mathcal{O}_M(\mathbb{R})$. Then $\varphi = C_\varphi p_1 \in \mathcal{O}_M(\mathbb{R})$. So, (3) is satisfied.

(2) \Leftrightarrow (3). Follows by Theorem 2.3.

(2) \Rightarrow (1). If the composition operator C_φ acts continuously from $\mathcal{O}^m(\mathbb{R})$ into itself for all $m \in \mathbb{N}$, taking into account of the fact that $\mathcal{O}_M(\mathbb{R})$ is the projective limit of the (LB)-spaces $\mathcal{O}^m(\mathbb{R})$, it follows that the operator C_φ acts continuously from $\mathcal{O}_M(\mathbb{R})$ into itself. \square

Remark 2.5. Let $\varphi \in \mathcal{O}_M(\mathbb{R})$. If φ is not a constant function, then the continuous linear operator $C_\varphi : \mathcal{O}_M(\mathbb{R}) \rightarrow \mathcal{O}_M(\mathbb{R})$ is never compact. The result follows from [13, Theorem 3.3], after having observed that $\mathcal{D}(\mathbb{R}) \hookrightarrow \mathcal{O}_M(\mathbb{R}) \hookrightarrow C^\infty(\mathbb{R})$ continuously.

Remark 2.6. Let $\mathcal{O}_C(\mathbb{R}) := \cup_{n=1}^{\infty} \cap_{m=1}^{\infty} \mathcal{O}_n^m(\mathbb{R})$ be the space of *very slowly increasing functions* on \mathbb{R} , endowed with its natural lc-topology, i.e., $\mathcal{O}_C(\mathbb{R}) = \text{ind}_{\rightarrow} \text{proj}_{\leftarrow} \mathcal{O}_n^m(\mathbb{R})$, which is a complete (LF)-space. In particular, $\mathcal{O}_C(\mathbb{R})$ is continuously included in $\mathcal{O}_M(\mathbb{R})$ and it is the predual of the space of convolutors of the spaces $\mathcal{S}(\mathbb{R})$ and $\mathcal{S}'(\mathbb{R})$.

Let $\varphi \in C^\infty(\mathbb{R})$. If C_φ acts continuously on $\mathcal{O}_C(\mathbb{R})$, then $\varphi \in \mathcal{O}_C(\mathbb{R})$ since $\varphi = C_\varphi p_1$, where $p_1 \in \mathcal{O}_C(\mathbb{R})$ is the identity function. But the converse is not true. For example, let $\varphi(x) = x^2$ and $f(x) = \sin x$, for $x \in \mathbb{R}$. Then $\varphi, f \in \mathcal{O}_C(\mathbb{R})$. But $f \circ \varphi \notin \mathcal{O}_C(\mathbb{R})$. Indeed, we have for every $x \in \mathbb{R}$ and $i \in \mathbb{N}$, with i even, that

$$(f \circ \varphi)^{(i)}(x) = P(x) \sin x^2 + Q(x) \cos x^2,$$

where P is a polynomial of degree equal to i and Q is a polynomial of degree less than i . So, $f \circ \varphi \notin \cap_{m=1}^{\infty} \mathcal{O}_n^m(\mathbb{R})$ for all $n \in \mathbb{N}$. Otherwise, if $f \circ \varphi \in \cap_{m=1}^{\infty} \mathcal{O}_{n_0}^m(\mathbb{R})$ for some $n_0 \in \mathbb{N}$, then

$$|(f \circ \varphi)|_{m, n_0} = \sup_{x \in \mathbb{R}} \sup_{j=0, \dots, m} (1+x^2)^{-n_0} |(f \circ \varphi)^j(x)| < +\infty$$

for all $m \in \mathbb{N}$. But if we choose $x_k^2 := \frac{\pi}{2} + 2k\pi$, for all $k \in \mathbb{N}$ and $i := 2(n_0 + 1)$, we get for all $k \in \mathbb{N}$ that

$$(1+x_k^2)^{-n_0} |(f \circ \varphi)^{2(n_0+1)}(x)| = (1+x_k^2)^{-n_0} |P(x_k)| \geq cx_k^2$$

for some suitable positive constant c . Therefore, $|(f \circ \varphi)|_{2(n_0+1), n_0} = +\infty$. A contradiction.

3. Dynamics of composition operators acting on lchS' continuously included in $C^m(\mathbb{R})$ for some $m \in \mathbb{N}_0 \cup \{\infty\}$

Through this section X always denotes a separable lchS. Let $T \in \mathcal{L}(X)$. The operator T is said to be *supercyclic* if there exists $x \in X$ whose projective orbit under T is dense in X , that is the set $\{\lambda T^n x : n \in \mathbb{N}_0, \lambda \in \mathbb{C}\}$ is dense in X . Such an x is said to be a *supercyclic vector* of T . The operator T is said to be *hypercyclic* if there exists $x \in X$ whose orbit under T is dense in X , that is the set $\text{orb}(x, T) := \{T^n x : n \in \mathbb{N}_0\}$ is dense in X . Such an x is said to be a *hypercyclic vector* of T . No power bounded operator is hypercyclic, but can be supercyclic. Every hypercyclic operator is always supercyclic. The converse is not true in general. The operator T is said to be *cyclic* if there exists $x \in X$ whose span of the orbit under T is dense in X , that is the space $\text{span}(\text{orb}(x, T))$ is dense in X . Such an x is said to be a *cyclic vector* of T . Every supercyclic operator is always cyclic. The converse is not true in general. The operator T is said to be *topological transitive* if for every pair of non-empty open subsets U and V of X , there is $n \in \mathbb{N}$ such that $T^n(U) \cap V \neq \emptyset$. An hypercyclic operator is always topological transitive. Thanks to the Birkhoff-transitivity theorem, the converse holds for Fréchet spaces (see [5, Theorem 1.2]).

We recall an important result that we use in the following. For the proof see [5, Proposition 1.26].

Proposition 3.1. *Let X be a separable lchS and let $T \in \mathcal{L}(X)$ be supercyclic. Then either $\sigma_p(T') = \emptyset$ or $\sigma_p(T') = \{\lambda\}$, for some $\lambda \neq 0$. In the latter case, $\text{Ker}(T' - \lambda)$ has dimension 1 and $\text{Ker}(T' - \lambda)^n = \text{Ker}(T' - \lambda)$ for all $n \in \mathbb{N}_0$. Moreover, there exists a (closed) T -invariant hyperplane $X_0 \subset X$ such that $T_0 := \lambda^{-1}T|_{X_0}$ is hypercyclic on X_0 .*

An operator $T \in \mathcal{L}(X)$ is said to be *mixing* if for every pair of non-empty open subsets U and V of X , there is $N \in \mathbb{N}$ such that $T^n(U) \cap V \neq \emptyset$ for all $n \geq N$. Every mixing operator on a separable Fréchet space X is hypercyclic. In particular, in such a case it is easy to see that if the operator T is mixing, then it is also *hereditarily hypercyclic*, i.e., for any infinite set $I \subseteq \mathbb{N}$ the family $\{T^n : n \in I\}$ is *universal* which means that for some $x \in X$ the set $\{T^n x : n \in I\}$ is dense in X .

An operator T is said to be *weakly hypercyclic* if there exists $x \in X$ whose orbit under T is weakly dense in X (for the weak topology $\sigma(X, X')$). Such a vector x is called a *weakly hypercyclic vector* for T . One defines in the same way *weakly supercyclic operators* and *weakly supercyclic vectors*. We refer the reader to [5,14] for more details.

In the following, if X is a lchS of functions defined on \mathbb{R} , we denote by δ_a the linear functional on X which maps every function of X to its value at $a \in \mathbb{R}$. We observe that if X is continuously included in $C^m(\mathbb{R})$, for some $m \in \mathbb{N}_0 \cup \{\infty\}$, then $\delta_a^{(j)} \in X'$ for all $j \leq m$ and $a \in \mathbb{R}$, where $\delta_a^{(j)}$ is defined by $\delta_a^{(j)}(f) := f^{(j)}(a)$. A function $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ is said to be *strongly runaway* when for any compact subset K of \mathbb{R} there exists $n_0 \in \mathbb{N}$ such that $\varphi_n(K) \cap K = \emptyset$ for all $n \geq n_0$. It is standard to show that for an increasing function $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ the property to be strongly runaway is equivalent to the fact that φ has no fixed points (cf. [20, Lemma 4.1] for a proof in the smooth case).

We now can prove the following result.

Proposition 3.2. *Let X be a separable lchS such that $X \hookrightarrow C^1(\mathbb{R})$ and $\delta_a, \delta_a^{(1)}$ are linearly independent in X' for every $a \in \mathbb{R}$. Let $\varphi \in C^1(\mathbb{R})$ be a symbol for X . If $\varphi(a) = a$ for some $a \in \mathbb{R}$ or $\varphi'(a) = 0$ for some $a \in \mathbb{R}$, then $C_\varphi : X \rightarrow X$ is not weakly supercyclic.*

Proof. Since φ is a symbol for X , $C_\varphi \in \mathcal{L}(X)$ and hence $C_\varphi \in \mathcal{L}(X, \sigma(X, X'))$.

Suppose firstly that $\varphi'(a) = 0$ for some $a \in \mathbb{R}$. This assumption implies that $C_\varphi(X) \subseteq \text{Ker } \delta_a^{(1)}$. Then C_φ cannot be even (weakly) cyclic.

Now, suppose that $\varphi(a) = a$ for some $a \in \mathbb{R}$. This assumption implies that $C'_\varphi(\delta_a) = \delta_a$ and $C'_\varphi(\delta_a^{(1)}) = \varphi'(a)\delta_{\varphi(a)}^{(1)} = \varphi'(a)\delta_a^{(1)}$. Now we have to distinguish two cases. If $\varphi'(a) \neq 1$, then C'_φ has two different eigenvalues. Applying Proposition 3.1 to $(X, \sigma(X, X'))$, we get that C_φ is not weakly supercyclic. If $\varphi'(a) = 1$, then δ_a and $\delta_a^{(1)}$ are two different eigenvectors for C'_φ , thereby implying that $\text{Ker}(C'_\varphi - I)$ has dimension greater or equal than two. Again, applying Proposition 3.1 to $(X, \sigma(X, X'))$, we get that C_φ is not weakly supercyclic. \square

Corollary 3.3. *Let X be a separable lchS such that $X \hookrightarrow C^1(\mathbb{R})$ and $\delta_a, \delta_a^{(1)}$ are linearly independent in X' for every $a \in \mathbb{R}$. Let $\varphi \in C^1(\mathbb{R})$ be a symbol for X . If C_φ is weakly supercyclic on X , then φ is strongly runaway and $\varphi'(x) > 0$ for all $x \in \mathbb{R}$.*

Proof. By Proposition 3.2 we can assume that $\varphi'(x) \neq 0$ for every $x \in \mathbb{R}$. Since Proposition 3.2 also implies that the function φ cannot have fixed points, it follows that φ is increasing (φ decreasing would have fixed points). Accordingly, the function $\varphi(x) - x$ has constant sign. From this it follows that all the orbits of φ are divergent. Hence φ is strongly runaway. \square

Observe that in Proposition 3.2, we ask that $X \hookrightarrow C^1(\mathbb{R})$. We now prove the following result which concerns the weaker assumption $X \hookrightarrow C(\mathbb{R})$.

Proposition 3.4. *Let X be a separable lchS such that $X \hookrightarrow C(\mathbb{R})$ with dense inclusion and $\{\delta_a : a \in \mathbb{R}\}$ is a linearly independent subset in X' . Let $\varphi \in C(\mathbb{R})$ be a symbol for X . If C_φ is supercyclic on X , then φ is increasing and without fixed points.*

Proof. The density of the inclusion $X \hookrightarrow C(\mathbb{R})$ permits us to consider only the case $X = C(\mathbb{R})$ in order to prove that C_φ supercyclic implies φ increasing and without fixed points.

If φ is not injective, then there exist $a, b \in \mathbb{R}$ with $a \neq b$ such that $\varphi(a) = \varphi(b)$. Then $C_\varphi(C(\mathbb{R})) \subseteq \text{Ker}(\delta_a - \delta_b)$, thereby implying that C_φ cannot be (even weakly) supercyclic.

We now suppose that φ has fixed points. If there are more than one, for instance $a, b \in \mathbb{R}$ with $a \neq b$, we have that $C'_\varphi(\delta_a) = \delta_a$ and $C'_\varphi(\delta_b) = \delta_b$. By Proposition 3.1 it follows that C_φ cannot be (even weakly) supercyclic.

Finally, we assume that φ has a unique fixed point, that without loss of generality we can suppose to be 0. Since φ is injective, the function $\varphi(x) - x$ has constant sign in $(-\infty, 0)$ and in $(0, \infty)$. Suppose φ to be increasing. If $\varphi(x) < x$ for all $x > 0$, then $\varphi_n(x) \rightarrow 0$ as $n \rightarrow \infty$ for all $x > 0$ and so C_φ is not (even weakly) supercyclic by [8, Theorem 8(ii)]. Hence, $\varphi(x) > x$ for every $x > 0$. In a similar way, we can show that $\varphi(x) < x$ for every $x < 0$. Therefore, the function φ is also surjective and $\varphi^{-1}(x) < x$ for every $x > 0$ and $\varphi^{-1}(x) > x$ for every $x < 0$. Accordingly, $\varphi^{-1} \in C(\mathbb{R})$. Since $C_\varphi : C(\mathbb{R}) \rightarrow C(\mathbb{R})$ is supercyclic, also $C_{\varphi^{-1}} : C(\mathbb{R}) \rightarrow C(\mathbb{R})$ is supercyclic. Again we get a contradiction by [8, Theorem 8(ii)]. If φ is not increasing, it must be decreasing since it is injective. The contradiction now follows taking into account of the fact that $C_\varphi : C(\mathbb{R}) \rightarrow C(\mathbb{R})$ is supercyclic if, and only if, $C_{\varphi_2} : C(\mathbb{R}) \rightarrow C(\mathbb{R})$ is supercyclic. \square

Combining our results with previous works of Bonet and Domański and Kalmes, we get the following result for dynamics of some composition operators. The proof of (i) below uses an unpublished argument due to Bonet and Domański which ensures that C_φ is sequentially supercyclic in $\mathcal{A}(\mathbb{R})$ for every analytic diffeomorphism φ in \mathbb{R} without fixed points.

Theorem 3.5.

- (i) *If a composition operator C_φ defined on $\mathcal{A}(\mathbb{R})$ is weakly supercyclic, then C_φ is mixing. If we assume in addition φ to be surjective, then C_φ is weakly supercyclic if, and only if, C_φ is sequentially hypercyclic.*
- (ii) *A composition operator C_φ defined on $C^m(\mathbb{R})$ with $m \in \mathbb{N} \cup \{\infty\}$ is weakly supercyclic if, and only if, C_φ is mixing.*
- (iii) *A composition operator C_φ defined on $C(\mathbb{R})$ is supercyclic if, and only if, C_φ is mixing.*

Proof. (i). By Corollary 3.3, if C_φ is weakly supercyclic then φ must be runaway, $\varphi'(x) \neq 0$ for all $x \in \mathbb{R}$, and φ does not have any fixed point. By [10, Theorem 2.3] C_φ is then mixing. If φ is also surjective, then φ is an analytic diffeomorphism, and therefore φ is analytically conjugate to $\psi(x) = x + 1$ by [6, Theorem 3.1]. We conclude since C_ψ is sequentially hypercyclic by [10, Theorem 3.6].

(ii) It follows by combining Corollary 3.3 with [18, Corollary 4.2].

Finally, (iii) is a consequence of Proposition 3.4 and [18, Corollary 4.1]. \square

Proposition 3.6. *Let $\varphi(x) = x + d$, $d \neq 0$. The operator C_φ is mixing on $\mathcal{O}^m(\mathbb{R})$ for any $m \in \mathbb{N}$, and also on $\mathcal{O}_M(\mathbb{R})$.*

Proof. Using Remark 2.2 it is easy to show that, for every $f \in \mathcal{D}(\mathbb{R})$, both sequences $\{f \circ \varphi_n\}_{n \in \mathbb{N}}$ and $\{f \circ \varphi_{-n}\}_{n \in \mathbb{N}}$ are convergent to 0 in $\mathcal{O}^m(\mathbb{R})$ for any $m \in \mathbb{N}$ and hence, in $\mathcal{O}_M(\mathbb{R})$. Now, by applying [5, Theorem 1.6] with $n_k = k$, $\mathcal{D}_1 = \mathcal{D}_2 = \mathcal{D}(\mathbb{R})$ and $S = C_\varphi$, we get that C_φ is mixing in $\mathcal{O}^m(\mathbb{R})$ for any $m \in \mathbb{N}$ and also in $\mathcal{O}_M(\mathbb{R})$. \square

The basic example of the translations gives us mixing operators on both spaces $\mathcal{O}^m(\mathbb{R})$, $m \in \mathbb{N}$, and $\mathcal{O}_M(\mathbb{R})$. This should be compared with [12, Theorem 2.1] and [12, Corollary 2.2(1)], which shows how the situation differs when $\{f \circ \varphi_n\}_{n \in \mathbb{N}}$ is not convergent to 0 when $f \in \mathcal{D}(\mathbb{R})$ and $\varphi(x) = x + d$. We do not know if $C_\varphi : \mathcal{O}_M(\mathbb{R}) \rightarrow \mathcal{O}_M(\mathbb{R})$ or $C_\varphi : \mathcal{O}^m(\mathbb{R}) \rightarrow \mathcal{O}^m(\mathbb{R})$ is mixing when φ is strongly runaway and $\varphi'(x) > 0$ for all $x \in \mathbb{R}$.

4. Ergodic properties of composition operators acting on lcHs' continuously included in $C^m(\mathbb{R})$ for some $m \in \mathbb{N}_0 \cup \{\infty\}$

4.1. Strong operator topology on operators on spaces of differentiable functions

We start by proving that the superposition operator is continuous.

Proposition 4.1. *Let $m \in \mathbb{N}_0 \cup \{\infty\}$ and let $f \in C^m(\mathbb{R})$. Then the superposition operator $S_f : C^m(\mathbb{R}) \rightarrow C^m(\mathbb{R})$, $\varphi \mapsto f \circ \varphi$, is continuous.*

Proof. We prove the statement by induction on m . Let $f \in C(\mathbb{R})$ and let $\{\varphi_n\}_{n \in \mathbb{N}} \subset C(\mathbb{R})$ be a convergent sequence in $C(\mathbb{R})$ to some $\varphi \in C(\mathbb{R})$. We will show that the sequence $\{f \circ \varphi_n\}_{n \in \mathbb{N}}$ is convergent to $f \circ \varphi$ in the topology τ_c of $C(\mathbb{R})$. To do this, we first observe that for all $k \in \mathbb{N}$ there exists $M \in \mathbb{N}$ such that $\varphi_n([-k, k]) \subseteq [-M, M]$ for all $n \in \mathbb{N}$. Indeed, there exists $n_0 \in \mathbb{N}$ such that $n \geq n_0$ implies $|\varphi_n(x) - \varphi(x)| \leq 1$ for each $x \in [-k, k]$. On the other hand, $\varphi([-k, k])$ and $\varphi_i([-k, k])$, $i = 1, \dots, n_0 - 1$, are compact subsets of \mathbb{R} as $\varphi, \varphi_i \in C(\mathbb{R})$, $i = 1, \dots, n_0 - 1$, and hence, $\varphi([-k, k]) \subset [-M', M']$ and $\varphi_i([-k, k]) \subset [-M', M']$, $i = 1, \dots, n_0 - 1$, for some $M' > 0$. Therefore, $\varphi_n([-k, k]) \subset [-M' - 1, M' + 1]$ for all $n \in \mathbb{N}$.

Since $f \in C(\mathbb{R})$, f is uniformly continuous in $[-M, M]$ and hence, for a fixed $\varepsilon > 0$, there exists $\delta > 0$ such that $|f(x) - f(y)| < \varepsilon$ for each $x, y \in [-M, M]$ with $|x - y| < \delta$. On the other hand, $\sup_{x \in [-k, k]} |\varphi_n(x) - \varphi(x)| \rightarrow 0$ for $n \rightarrow \infty$ and hence, there exists $n_0 \in \mathbb{N}$ such that $\sup_{x \in [-k, k]} |\varphi_n(x) - \varphi(x)| < \delta$ for all $n \geq n_0$. Since $\varphi_n([-k, k]) \subset [-M, M]$ for all $n \in \mathbb{N}$, we get that $|(f \circ \varphi_n)(x) - (f \circ \varphi)(x)| < \varepsilon$ for all $n \geq n_0$ and $x \in [-k, k]$.

Assume now the hypothesis is true for $m - 1$. Let $f \in C^m(\mathbb{R})$ and let $\{\varphi_n\}_{n \in \mathbb{N}} \subset C^m(\mathbb{R})$ be a convergent sequence in $C^m(\mathbb{R})$ to some $\varphi \in C^m(\mathbb{R})$. For all $n \in \mathbb{N}$, for every $x \in \mathbb{R}$ and $i = 0, \dots, m$, Faà di Bruno formula gives:

$$(S_f(\varphi_n))^{(i)}(x) = \sum \frac{i!}{k_1!k_2! \dots k_n!} f^{(k)}(\varphi_n(x)) \left(\frac{\varphi_n'(x)}{1!}\right)^{k_1} \left(\frac{\varphi_n''(x)}{2!}\right)^{k_2} \dots \left(\frac{\varphi_n^{(i)}(x)}{i!}\right)^{k_i}, \tag{4.1}$$

where the sum is extended over all $(k_1, k_2, \dots, k_i) \in \mathbb{N}_0^i$ such that $k_1 + 2k_2 + \dots + ik_i = i$ and $k_1 + k_2 + \dots + k_i = k$. We get from (4.1), the inductive hypothesis and the continuity of the product in $C(\mathbb{R})$ that $\{(S_f(\varphi_n))^{(i)}\}_{n \in \mathbb{N}}$ is convergent to $(S_f(\varphi))^{(i)}$ in $C(\mathbb{R})$ for any $0 \leq i \leq m$, thereby completing the proof. \square

Corollary 4.2. *Let $m \in \mathbb{N}_0 \cup \{\infty\}$ and X be a barreled lcHs continuously included in $C^m(\mathbb{R})$ and satisfying the following properties:*

- (i) *The identity $p_1(x) = x \in X$;*
- (ii) *X is closed under composition;*
- (iii) *A sequence $\{\varphi_n\}_{n \in \mathbb{N}} \subset X$ is convergent in X to some $\varphi \in X$ if, and only if, $\{\varphi_n\}_{n \in \mathbb{N}}$ is bounded in X and $\lim_{n \rightarrow \infty} \varphi_n = \varphi$ in $C^m(\mathbb{R})$.*

Let $\{\varphi_n\}_{n \in \mathbb{N}} \subset X$ be a sequence of symbols for X . Then the sequence $\{C_{\varphi_n}\}_{n \in \mathbb{N}}$ of composition operators is convergent in $\mathcal{L}_s(X)$ if, and only if, $\{\varphi_n\}_{n \in \mathbb{N}}$ is convergent in X and $\{C_{\varphi_n}\}_{n \in \mathbb{N}}$ is an equicontinuous sequence in $\mathcal{L}(X)$.

Proof. If there exists $C \in \mathcal{L}(X)$ such that $\{C_{\varphi_n}(f)\}_{n \in \mathbb{N}}$ is convergent to $C(f)$ for any $f \in X$, then for $f = p_1$ we get that $\{\varphi_n\}_{n \in \mathbb{N}}$ converges in X to $C(p_1)$. Since X is barreled, we also obtain that $\{C_{\varphi_n}\}_{n \in \mathbb{N}}$ is an equicontinuous sequence in $\mathcal{L}(X)$.

The converse follows from Proposition 4.1. Indeed, if the sequence $\{\varphi_n\}_{n \in \mathbb{N}}$ is convergent in X to some $\varphi \in X$, then by (iii) the sequence $\{\varphi_n\}_{n \in \mathbb{N}}$ is convergent in $C^m(\mathbb{R})$ to φ and so, by Proposition 4.1 we have that $C_{\varphi_n}(f) = f \circ \varphi_n \rightarrow f \circ \varphi$ in $C^m(\mathbb{R})$ as $n \rightarrow \infty$ for any fixed $f \in X$. On the other hand, the sequence $\{C_{\varphi_n}(f)\}_{n \in \mathbb{N}}$ is bounded in X for any fixed $f \in X$, being $\{C_{\varphi_n}\}_{n \in \mathbb{N}}$ an equicontinuous sequence in $\mathcal{L}(X)$. Therefore, by (iii) we conclude that $C_{\varphi_n}(f) = f \circ \varphi_n \rightarrow f \circ \varphi$ in X as $n \rightarrow \infty$ for any fixed $f \in X$. The equicontinuity of $\{C_{\varphi_n}\}_{n \in \mathbb{N}}$ implies that $C_\varphi \in \mathcal{L}(X)$. So, $C_{\varphi_n} \rightarrow C_\varphi$ in $\mathcal{L}_s(X)$ as $n \rightarrow \infty$. \square

Remark 4.3. If X is a Montel lchS space which is continuously included in $C^m(\mathbb{R})$, with $m \in \mathbb{N}_0 \cup \{\infty\}$, then X clearly satisfies condition (iii) of Corollary 4.2.

Remark 4.4. If the sequence $\{\varphi_n\}_{n \in \mathbb{N}}$ is bounded in $C^m(\mathbb{R})$ for some $m \in \mathbb{N}_0 \cup \{\infty\}$, then for all $k \in \mathbb{N}$ there exists $M > 0$ such that $\sup_{|x| \leq k} |\varphi_n^{(j)}(x)| < M$ for all $0 \leq j \leq m$ and $n \in \mathbb{N}$, i.e., $\varphi_n^{(j)}([-k, k]) \subseteq [-M, M]$ for all $0 \leq j \leq m$ and $n \in \mathbb{N}$. Hence, [17, Theorem 3.13] for the particular case of composition operators reads as follows: C_φ is power bounded in $C^m(\mathbb{R})$ if, and only if, $\{\varphi_n\}_{n \in \mathbb{N}}$ is bounded in $C^m(\mathbb{R})$.

4.2. Polynomial symbols

We collect some results on mean ergodicity of C_φ .

Proposition 4.5. Let X be a lchS which is closed under composition, continuously included in $C^m(\mathbb{R})$ for some $m \in \mathbb{N}_0 \cup \{\infty\}$ and contains the function $p_1(x) := x$, for $x \in \mathbb{R}$. If $\varphi \in X$ is a symbol for X and C_φ is mean ergodic on X , then the following properties are satisfied:

- (1) The sequence $\{\varphi_{[n]}\}_{n \in \mathbb{N}}$ converges in $C^m(\mathbb{R})$;
- (2) The sequence $\{\frac{\varphi_n}{n}\}_{n \in \mathbb{N}}$ converges to 0 in $C^m(\mathbb{R})$.

If C_φ is power bounded on X , condition (2) above is satisfied.

Proof. Assume that C_φ is mean ergodic on X .

(1) By assumption there exists $P \in X$ such that $(C_\varphi)_{[n]} \rightarrow P$ in $\mathcal{L}_s(X)$. Since the function p_1 belongs to X , it follows that $(C_\varphi)_{[n]}(p_1) = \varphi_{[n]} \rightarrow Pp_1 := \psi$ in X . Hence, $\{\varphi_{[n]}\}_{n \in \mathbb{N}}$ is also convergent in $C^m(\mathbb{R})$ to ψ . Therefore, condition (1) follows.

(2) Since

$$\frac{\varphi_n}{n} = \varphi_{[n]} - \frac{n-1}{n} \varphi_{[n-1]}, \quad n \geq 2,$$

the result immediately follows from part (1).

Finally, if C_φ is power bounded, then the sequence $\{C_{\varphi_n}(p_1)\}_{n \in \mathbb{N}} = \{\varphi_n\}_{n \in \mathbb{N}}$ is bounded in X and hence, bounded in $C^m(\mathbb{R})$, thereby implying that (2) is satisfied. \square

Theorem 4.6. Let φ be a polynomial and let $X \in \{\mathcal{O}_M(\mathbb{R}), C^\infty(\mathbb{R}), \mathcal{A}(\mathbb{R})\} \cup \{\mathcal{O}^m(\mathbb{R}) : m \in \mathbb{N}_0\} \cup \{C^m(\mathbb{R}) : m \in \mathbb{N}_0\}$. Then the composition operator C_φ is mean ergodic on X if, and only if, C_φ is power bounded if, and only if, $\varphi(x) = ax + b$, with either $|a| < 1$, $a = -1$ or $a = 1$ and $b = 0$. In case $|a| < 1$, we even have that the sequence of iterates $\{C_{\varphi_n}\}_{n \in \mathbb{N}}$ converges in $\mathcal{L}_s(X)$.

Proof. Assume that either the degree of φ is greater or equal than two or $\varphi(x) = ax + b$ with $|a| > 1$. Then there exists $n_0 \in \mathbb{N}$ such that $|x| \geq n_0$ implies $|\varphi(x)| > |x| + 1$. So, it follows that $|\varphi_n(n_0)| \geq n_0 + n$ for all $n \in \mathbb{N}$. Therefore, C_φ is neither power bounded nor mean ergodic on X by Proposition 4.5. If $\varphi(x) = x + b$,

with $b \neq 0$, then $\varphi_n(0) = nb$, and again Proposition 4.5 yields that C_φ cannot be power bounded or mean ergodic on X .

Let $\varphi(x) := ax + b$ for $x \in \mathbb{R}$ and some $a, b \in \mathbb{R}$. The case $a = 1, b = 0$ gives the identity. The case $a = -1$ gives $\varphi_2 = \varphi$, and hence, C_φ is idempotent. In both cases the statement is trivial. Also it is trivial when $a = 0$. Indeed, in this case C_φ is the evaluation at b and satisfies $C_{\varphi_n} = C_\varphi$ for all $n \in \mathbb{N}$.

Let $|a| < 1$, with $a \neq 0$. In this case, for $\phi(x) := x + b/(a - 1)$, and $\psi(x) := ax$, we have $C_\varphi = C_{\phi^{-1} \circ \psi \circ \phi} = C_\phi C_\psi C_{\phi^{-1}}$. Hence, power boundedness or mean ergodicity of C_φ is equivalent to that of C_ψ . Thus, we can assume $b = 0$. In this case it is easy to show that the sequence $\{\varphi_n\}_{n \in \mathbb{N}}$ is convergent to the constant function 0 in all the spaces we are considering and hence, bounded there. Accordingly, the operator C_φ is power bounded in every space X (see Theorem 5.2 for X equals to $\mathcal{O}_M(\mathbb{R})$ or equals to $\mathcal{O}^m(\mathbb{R})$ with $m \in \mathbb{N}$; see Remark 4.4 for X equals to $C^m(\mathbb{R})$ with $m \in \mathbb{N}_0 \cup \{\infty\}$; for $X = \mathcal{A}(\mathbb{R})$ see [9]). Since every space X satisfies the assumptions (i)–(iii) of Corollary 4.2, the conclusion follows from Corollary 4.2. \square

We point out that if φ is a polynomial of even degree and with no fixed points, like $\varphi(x) = x^2 + 1$ for $x \in \mathbb{R}$, then the composition operator C_φ is power bounded on $\mathcal{S}(\mathbb{R})$ (see [12, Theorem 3.7]). The key point to make the difference is that in $\mathcal{S}(\mathbb{R})$ the polynomials are symbols, but they do not belong to $\mathcal{S}(\mathbb{R})$. We see below that in case of analytic functions we can go further than polynomial symbols.

Remark 4.7. Bonet and Domanski showed in [9] that a nontrivial composition operator C_φ (i.e., $\varphi_2 \neq p_1$) defined on $\mathcal{A}(\mathbb{R})$ is power bounded if, and only if, $\{\varphi_n\}_{n \in \mathbb{N}}$ is convergent in $\mathcal{A}(\mathbb{R})$ to a (unique) fixed point a of φ . Hence, from Corollary 4.2 we get that such an operator satisfies that $\{C_{\varphi_n}\}_{n \in \mathbb{N}}$ is convergent in $\mathcal{L}_b(\mathcal{A}(\mathbb{R}))$.

4.3. Monotonic symbols

Proposition 4.8. *Let X be a lcHs which is closed under composition, continuously included in $C(\mathbb{R})$ and contains the function p_1 . Let $\varphi \in X$ be a symbol for X such that there exists $\lim_{n \rightarrow \infty} \varphi_n(x)$ for all $x \in \mathbb{R}$. If C_φ is mean ergodic on X , then the set $\text{Fix}(\varphi)$ of fixed points of φ is a non empty closed interval of \mathbb{R} .*

Proof. Since C_φ is mean ergodic on X , we can apply Proposition 4.5(1) to get that the sequence $\{\varphi_{[n]}\}_{n \in \mathbb{N}}$ converges in $C(\mathbb{R})$ to some function $\psi \in C(\mathbb{R})$. Since there exists $\lim_{n \rightarrow \infty} \varphi_n(x)$ for every $x \in \mathbb{R}$, by Cesàro’s theorem we necessary have

$$\psi(x) = \lim_{n \rightarrow \infty} \varphi_n(x)$$

for every $x \in \mathbb{R}$. So, by the fact that φ is a continuous function, we also have for every $x \in \mathbb{R}$ that

$$\psi(x) = \lim_{n \rightarrow \infty} \varphi_n(x) = \lim_{n \rightarrow \infty} \varphi(\varphi_{n-1}(x)) = \varphi(\psi(x)).$$

This means that $\psi(x) \in \text{Fix}(\varphi)$ for all $x \in \mathbb{R}$ and hence, $\text{Fix}(\varphi) = \text{Im}\psi$. Since $\text{Fix}(\varphi)$ is always a closed set and $\psi \in C(\mathbb{R})$, it follows that $\text{Fix}(\varphi)$ is a non empty closed interval of \mathbb{R} . \square

Proposition 4.9. *Let X be a lcHs which is closed under composition, continuously included in $C(\mathbb{R})$ and contains the function p_1 . Let $\varphi \in X$ be a symbol for X . If there exist $\beta > 0$ such that φ is a non decreasing function on $[\beta, +\infty)$ (on $(-\infty, \beta]$, respectively), and $\varphi(x) > x$ ($\varphi(x) < x$) for every $x \geq \beta$ ($x \leq \beta$, respectively), then C_φ is not mean ergodic on X .*

Proof. We consider only the case in which φ is a non decreasing function on $[\beta, +\infty)$. In the other case the proof follows by arguing in a similar way.

We first observe that $\{\varphi_n(\beta)\}_{n \in \mathbb{N}}$ is a non decreasing sequence, but not a bounded sequence. Indeed, from $\varphi(\beta) > \beta$ follows that there exists $\delta > 0$ such that $\varphi(\beta) > \beta + \delta$ and hence, $\varphi_n(\beta) > \beta + n\delta$ for all $n \in \mathbb{N}$. Accordingly, $\frac{\varphi_n(\beta)}{n} > \frac{\beta}{n} + \delta > \delta$ for all $n \in \mathbb{N}$. Hence, $\{\frac{\varphi_n(\beta)}{n}\}_{n \in \mathbb{N}}$ does not converge to 0. So, by Proposition 4.5 we can conclude that C_φ is not mean ergodic on X . \square

Corollary 4.10. *Let X be a lchS which is closed under composition, continuously included in $C^1(\mathbb{R})$ and contains the function p_1 . Let $\varphi \in X$ be a symbol for X , $\varphi \neq p_1$. If φ is a non decreasing function and C_φ is mean ergodic on X , then there is a $a \in \mathbb{R}$ such that $\text{Fix}(\varphi) = \{a\}$ and $\{\varphi_n\}_{n \in \mathbb{N}}$ converges in $C(\mathbb{R})$ to the constant function $\psi(x) := a$ for $x \in \mathbb{R}$.*

Proof. Since φ is a non decreasing function, there exists $\lim_{n \rightarrow \infty} \varphi_n(x)$ for all $x \in \mathbb{R}$. Indeed, for a fixed $x \in \mathbb{R}$, we have either $\varphi(x) \leq x$ or $\varphi(x) \geq x$ from which it follows that $\{\varphi_n(x)\}_{n \in \mathbb{N}}$ is either a non increasing sequence or a non decreasing sequence, being φ a non decreasing function. Since C_φ is mean ergodic, we can then apply Proposition 4.8 to conclude that the sequence $\{\varphi_{[n]}\}_{n \in \mathbb{N}}$ converges in X (and hence, in $C^1(\mathbb{R})$) to some $\psi \in X$ and the set $\text{Fix}(\varphi) = \psi(\mathbb{R})$ is a closed interval of \mathbb{R} . The pointwise convergence of $(\varphi_{[n]})_{n \in \mathbb{N}}$ to ψ together with the monotonicity of $(\varphi_n(x))_{n \in \mathbb{N}}$ for all $x \in \mathbb{R}$ implies that $(\varphi_n)_{n \in \mathbb{N}}$ is pointwise convergent and $\psi(x) = \lim_{n \rightarrow \infty} \varphi_n(x)$ for every $x \in \mathbb{R}$.

Let assume $\text{Fix}(\varphi)$ to be bounded below and let a be the infimum of $\text{Fix}(\varphi)$. Clearly, $a \in \text{Fix}(\varphi)$. On the other hand, since C_φ is mean ergodic, Proposition 4.9 implies that $\varphi(x) > x$ for all $x < a$. Accordingly, taking into account that φ is a non decreasing function, it follows for every $x < a$ that $\{\varphi_n(x)\}_{n \in \mathbb{N}}$ is a non decreasing sequence such that $\varphi_n(x) \leq \varphi_n(a) = a$ for all $n \in \mathbb{N}$. Hence, $\psi(x) = \lim_n \varphi_n(x) \leq a$ for all $x < a$. But, we also have for every $x < a$ that $\psi(x) = \lim_{n \rightarrow \infty} \varphi(\varphi_{n-1}(x)) = \varphi(\psi(x))$ and hence, $\psi(x) \in \text{Fix}(\varphi)$, thereby implying that $\psi(x) \geq a$. So, $\psi(x) = a$ for all $x < a$. If $\text{Fix}(\varphi)$ is not a single point, then there is $\delta > 0$ such that $[a, a + \delta) \subseteq \text{Fix}(\varphi)$. For all $x \in [a, a + \delta)$ we have $\psi(x) = \lim_n \varphi_{[n]}(x) = x$. This contradicts the differentiability of ψ at a .

The proof of the case $\text{Fix}(\varphi)$ bounded above is completely analogous.

Now, we can assume that $a = 0$ without loss of generality. So, Proposition 4.9 yields $|\varphi(x)| \leq |x|$ for all $x \in \mathbb{R}$. From this we get immediately that $\lim_{n \rightarrow \infty} \varphi_n(x) = 0$ uniformly on compact subsets of \mathbb{R} . \square

Theorem 4.11. *Let X be a Montel lchS which is closed under composition, continuously included in $C^1(\mathbb{R})$ and contains the function p_1 . Assume that the following condition is also satisfied:*

(*) *The composition operator $C_\varphi : X \rightarrow X$ is power bounded if, and only if, $\{\varphi_n\}_{n \in \mathbb{N}}$ is bounded in X for any $\varphi \in X$ symbol for X .*

Let $\varphi \in X$ be monotonic, $\varphi_2 \neq p_1$. Then C_φ is power bounded if, and only if, φ has an attracting fixed point a and $\{C_{\varphi_n}\}_{n \in \mathbb{N}}$ is convergent to C_a in $\mathcal{L}_s(X)$.

Proof. We first observe that if C_φ is power bounded, then the sequence $\{\varphi_n\}_{n \in \mathbb{N}}$ is bounded in X , being $\varphi_n = C_{\varphi_n} p_1$ for all $n \in \mathbb{N}$.

Assume first that φ is non decreasing. Since X is Montel and C_φ is power bounded, C_φ is also (uniformly) mean ergodic. Thus, by Corollary 4.10 the sequence $\{\varphi_n\}_{n \in \mathbb{N}}$ necessarily converges in $C(\mathbb{R})$ to a unique fixed point a . Now, the fact that X is Montel together with (*) imply that $\{\varphi_n\}_{n \in \mathbb{N}}$ is also convergent in X to a . Therefore, we can apply Corollary 4.2 to conclude the convergence of $\{C_{\varphi_n}\}_{n \in \mathbb{N}}$ to C_a in $\mathcal{L}_s(X)$, taking into account that C_φ is power bounded. Vice versa, if $\{C_{\varphi_n}\}_{n \in \mathbb{N}}$ is convergent in $\mathcal{L}_s(X)$ to C_a , then C_φ is power bounded, and we conclude.

Assume now that φ is decreasing. Since X is barreled, by Banach-Steinhaus theorem we have that any operator $T \in \mathcal{L}(X)$ is power bounded if, and only if, T^2 is. Accordingly, C_φ is power bounded if, and only

if, C_{φ_2} is. Since φ_2 is increasing, by the first part of the proof we get that C_φ is power bounded if, and only if, φ_2 has only one fixed point a and $\{C_{\varphi_{2n}}\}_{n \in \mathbb{N}}$ is convergent to C_a in $\mathcal{L}_s(X)$. It follows that also φ has only one fixed point a . On the other hand, the convergence of $\{C_{\varphi_{2n}}\}_{n \in \mathbb{N}}$ to C_a in $\mathcal{L}_s(X)$ implies the convergence of $\{\varphi_{2n}\}_{n \in \mathbb{N}}$ to the constant function a in $C(\mathbb{R})$. By applying this to any $\varphi([\alpha, \beta])$, with $\alpha, \beta \in \mathbb{R}$, we obtain the convergence of $\{\varphi_n\}_{n \in \mathbb{N}}$ to the constant function a in $C(\mathbb{R})$. Taking account of the fact that $\{\varphi_n\}_{n \in \mathbb{N}}$ is a bounded sequence of X as C_φ is power bounded, we can apply Corollary 4.2 to conclude that $\{C_{\varphi_n}\}_{n \in \mathbb{N}}$ is convergent to C_a in $\mathcal{L}_s(X)$. \square

5. Power boundedness of composition operators on $\mathcal{O}_M(\mathbb{R})$ and on $\mathcal{O}^m(\mathbb{R})$

A characterization of the power boundedness of C_φ when it acts continuously on $\mathcal{S}(\mathbb{R})$ has been proved in [12]. Precisely, in [12, Proposition 4] it has been shown that if $\varphi \in C^\infty(\mathbb{R})$ is a symbol for $\mathcal{S}(\mathbb{R})$, then the composition operator C_φ is power bounded if, and only if, the following conditions are satisfied:

- (1) For all $j \in \mathbb{N}_0$ there exist $C, p > 0$ such that, for every $x \in \mathbb{R}$ and $n \in \mathbb{N}$

$$|\varphi_n^{(j)}(x)| \leq C(1 + \varphi_n(x)^2)^p;$$

- (2) There exists $k > 0$ such that

$$|\varphi_n(x)| \geq |x|^{\frac{1}{k}}, \quad \forall |x| \geq k, \forall n \in \mathbb{N}.$$

In order to treat the case of the space $\mathcal{O}_M(\mathbb{R})$ we first establish the following result.

Theorem 5.1. *Let $m \in \mathbb{N}$ and $\varphi \in \mathcal{O}^m(\mathbb{R})$. Then the following properties are equivalent:*

- (1) C_φ is power bounded on $\mathcal{O}^m(\mathbb{R})$;
- (2) The sequence $\{\varphi_n\}_{n \in \mathbb{N}}$ is bounded in $\mathcal{O}^m(\mathbb{R})$;
- (3) For all $0 \leq i \leq m$ there exist $C, p > 0$ such that $|\varphi_n^{(i)}(x)| \leq C(1 + x^2)^p$ for all $x \in \mathbb{R}$ and $n \in \mathbb{N}$.

Proof. (1) \Rightarrow (2). The operator C_φ is power bounded on $\mathcal{O}^m(\mathbb{R})$. Accordingly, if we consider the function $p_1(x) := x$, for $x \in \mathbb{R}$, which belongs to $\mathcal{O}^m(\mathbb{R})$, then by assumption the sequence $\{C_\varphi^n p_1\}_{n \in \mathbb{N}} = \{\varphi_n\}_{n \in \mathbb{N}}$ is bounded in $\mathcal{O}^m(\mathbb{R})$.

(2) \Rightarrow (1). We claim that for all $r \in \mathbb{N}$ there exists $r' \in \mathbb{N}$ such that $C_\varphi: \mathcal{O}_{r'}^m(\mathbb{R}) \rightarrow \mathcal{O}_r^m(\mathbb{R})$ is power bounded, thereby obtaining that the sequence $\{f \circ \varphi_n\}_{n \in \mathbb{N}}$ is bounded in $\mathcal{O}_{r'}^m(\mathbb{R})$ for all $f \in \mathcal{O}_r^m(\mathbb{R})$. Indeed, for fixed $r \in \mathbb{N}$ and $f \in \mathcal{O}_r^m(\mathbb{R})$, we observe for every $x \in \mathbb{R}$ and $n \in \mathbb{N}$ that

$$(C_\varphi^n f)^{(i)}(x) = \sum \frac{i!}{k_1! k_2! \dots k_i!} f^{(k)}(\varphi_n(x)) \left(\frac{\varphi_n'(x)}{1!}\right)^{k_1} \left(\frac{\varphi_n''(x)}{2!}\right)^{k_2} \dots \left(\frac{\varphi_n^{(i)}(x)}{i!}\right)^{k_i},$$

where the sum is extended over all $(k_1, k_2, \dots, k_i) \in \mathbb{N}_0^i$ such that $k_1 + 2k_2 + \dots + ik_i = i$, with $k = k_1 + \dots + k_i$. Since by assumption on $\{\varphi_n\}_{n \in \mathbb{N}}$ (i.e., $\{\varphi_n\}_{n \in \mathbb{N}}$ is a bounded sequence of $\mathcal{O}^m(\mathbb{R})$), there exist $D > 0$ and $p \in \mathbb{N}$ such that $|\varphi_n^{(j)}(x)| \leq D(1 + x^2)^p$ for every $x \in \mathbb{R}$, $j = 0, \dots, m$ and $n \in \mathbb{N}$, and hence, $(1 + \varphi_n(x)^2) \leq D'(1 + x^2)^{2p}$ for every $x \in \mathbb{R}$ and $n \in \mathbb{N}$, and a suitable constant $D' > 0$, it follows for every $x \in \mathbb{R}$, $i = 0, \dots, m$ and $n \in \mathbb{N}$ that

$$|(C_\varphi^n f)^{(i)}(x)| \leq \sum \frac{i!}{k_1! k_2! \dots k_i!} |f^{(k)}(\varphi_n(x))| \left|\frac{\varphi_n'(x)}{1!}\right|^{k_1} \left|\frac{\varphi_n''(x)}{2!}\right|^{k_2} \dots \left|\frac{\varphi_n^{(i)}(x)}{i!}\right|^{k_i}$$

$$\begin{aligned} &\leq \sum \frac{i!}{k_1!k_2!\dots k_i!} |f|_{m,r} (1 + \varphi_n(x)^2)^r \frac{D^k(1+x^2)^{pk}}{1!k_1 2!k_2 \dots i!k_i} \\ &\leq C_i |f|_{m,r} (1+x^2)^{p(2r+i)} \leq C |f|_{m,r} (1+x^2)^{p(2r+m)}, \end{aligned}$$

where $C_i := (D')^r \sum \frac{i!}{k_1!k_2!\dots k_i!} \frac{D^k}{1!k_1 2!k_2 \dots i!k_i}$ and $C := \max_{i=0,\dots,m} C_i$. Accordingly, if $r' := p(2r+m) \in \mathbb{N}$, we get for all $n \in \mathbb{N}$ that

$$|C_\varphi^n f|_{m,r'} \leq C |f|_{m,r}.$$

Since $r \in \mathbb{N}$ and $f \in \mathcal{O}_r^m(\mathbb{R})$ are arbitrary, the claim is proved.

We are now able to show that C_φ is power bounded. Indeed, for a fixed $f \in \mathcal{O}^m(\mathbb{R})$, there exists $r \in \mathbb{N}$ such that $f \in \mathcal{O}_r^m(\mathbb{R})$. By the proof above, it follows that the sequence $\{C_\varphi^n f\}_{n \in \mathbb{N}}$ is bounded in $\mathcal{O}_{r'}^m(\mathbb{R})$ for some $r' \geq r$ and hence in $\mathcal{O}^m(\mathbb{R}) = \text{ind}_h \mathcal{O}_h^m(\mathbb{R})$, being it an (LB)-space. So, we can conclude that C_φ is power bounded on $\mathcal{O}^m(\mathbb{R})$ thanks to the Banach-Steinhaus theorem.

(2) \Leftrightarrow (3). Follows as $\mathcal{O}^m(\mathbb{R})$ is a regular (LB)-space. \square

Thanks to Theorem 5.1, we easily obtain the following characterization of the power boundedness of C_φ when it acts in $\mathcal{O}_M(\mathbb{R})$.

Theorem 5.2. *Let $\varphi \in \mathcal{O}_M(\mathbb{R})$. Then the following properties are equivalent:*

- (1) C_φ is power bounded on $\mathcal{O}_M(\mathbb{R})$;
- (2) C_φ is power bounded on $\mathcal{O}^m(\mathbb{R})$ for all $m \in \mathbb{N}$;
- (3) The sequence $\{\varphi_n\}_{n \in \mathbb{N}}$ is bounded in $\mathcal{O}_M(\mathbb{R})$;
- (4) For all $i \in \mathbb{N}_0$ there exist $C, p > 0$ such that $|\varphi_n^{(i)}(x)| \leq C(1+x^2)^p$ for all $x \in \mathbb{R}$ and $n \in \mathbb{N}$.

Proof. Since $\mathcal{O}_M(\mathbb{R})$ is the projective limit of the sequence $(\mathcal{O}^m(\mathbb{R}))_{m \in \mathbb{N}}$ of (LB)-spaces and it is barreled, the Banach-Steinhaus theorem yields that C_φ is power bounded if, and only if, $\{f \circ \varphi_n\}_{n \in \mathbb{N}}$ is bounded in $\mathcal{O}^m(\mathbb{R})$ for any $m \in \mathbb{N}$ and any $f \in \mathcal{O}_M(\mathbb{R})$. So, the conclusion follows now immediately from Theorem 5.1. \square

From Remark 4.4, Theorems 5.1 and 5.2 we get immediately the following fact.

Proposition 5.3. *Let $\varphi \in \mathcal{O}_M(\mathbb{R})$ ($\varphi \in \mathcal{O}^m(\mathbb{R})$, for $m \in \mathbb{N}$, resp.). Then C_φ is power bounded in $C^\infty(\mathbb{R})$ (in $\mathcal{C}^m(\mathbb{R})$, resp.) whenever C_φ is power bounded in $\mathcal{O}_M(\mathbb{R})$ (in $\mathcal{O}^m(\mathbb{R})$, resp.).*

Combining Theorem 4.11 with Theorem 5.2, we immediately obtain the following characterization, that should be compared with [17, Theorem 3.13].

Corollary 5.4. *Let $\varphi \in \mathcal{O}_M(\mathbb{R})$ be an increasing function. Then $\{C_{\varphi_n}\}_{n \in \mathbb{N}}$ is convergent in $\mathcal{L}_s(\mathcal{O}_M(\mathbb{R}))$ if, and only if, C_φ is power bounded on $\mathcal{O}_M(\mathbb{R})$ if, and only if, $\{\varphi_n\}_{n \in \mathbb{N}}$ is bounded in $\mathcal{O}_M(\mathbb{R})$.*

Acknowledgments

This paper was completed during the visit of the third author to the Universitat Politècnica de València. He expresses his gratitude to José Bonet and his collaborators for their kind hospitality and for their valuable suggestions and comments.

The research of E. Jordá was partially supported by the project PID2020-119457GB-100 funded by MCIN / AEI / 10.13039 / 501100011033 and by “ERDF A way of making Europe”. The research of E. Jordá is also partially supported by GVA-AICO/2021/170.

The authors would like to thank the referee for his/her careful reading of the work and his/her valuable suggestions, which certainly improved the work.

References

- [1] A.A. Albanese, J. Bonet, W.J. Ricker, Mean ergodic operators in Fréchet spaces, *Ann. Acad. Sci. Fenn., Math.* 34 (2009) 401–436.
- [2] A.A. Albanese, J. Bonet, W.J. Ricker, Convergence of arithmetic means of operators in Fréchet spaces, *J. Math. Anal. Appl.* 401 (2013) 160–173.
- [3] A.A. Albanese, J. Bonet, W.J. Ricker, Uniform convergence and spectra of operators in a class of Fréchet spaces, *Abstr. Appl. Anal.* 2014 (2014) 179027.
- [4] A.A. Albanese, C. Mele, Spectra and ergodic properties of multiplication and convolution operators on the space $\mathcal{S}(\mathbb{R})$, *Rev. Mat. Complut.* (2021), <https://doi.org/10.1007/s13163-021-00403-0>.
- [5] F. Bayart, E. Matheron, *Dynamics of Linear Operators*, Cambridge Tracts in Mathematics, vol. 179, Cambridge University Press, Cambridge, 2009.
- [6] G. Belitskii, Y. Lyubich, The real-analytic solutions of the Abel functional equation, *Stud. Math.* 134 (2) (1999) 135–141.
- [7] M.J. Beltrán-Meneu, E. Jordá, Dynamics of weighted composition operators on spaces of entire functions of exponential and infraexponential type, *Mediterr. J. Math.* 18 (5) (2021) 212.
- [8] M.J. Beltrán-Meneu, E. Jordá, M. Murillo-Arcila, Supercyclicity of weighted composition operators on spaces of continuous functions, *Collect. Math.* 71 (2020) 493–509.
- [9] J. Bonet, P. Domański, Power bounded composition operators on spaces of analytic functions, *Collect. Math.* 62 (2011) 69–83.
- [10] J. Bonet, P. Domański, Hypercyclic composition operators on spaces of real analytic functions, *Math. Proc. Camb. Philos. Soc.* 153 (2012) 489–503.
- [11] C. Chevalley, *Theory of Distributions: Lectures*, Columbia University, 1951.
- [12] C. Fernández, A. Galbis, E. Jordá, Dynamics and spectra of composition operators on the Schwartz space, *J. Funct. Anal.* 274 (2018) 3503–3530.
- [13] A. Galbis, E. Jordá, Composition operators on the Schwartz space, *Rev. Mat. Iberoam.* 34 (2018) 397–412.
- [14] K.G. Grosse-Erdmann, A. Peris, *Linear Chaos*, Springer, London, 2011.
- [15] A. Grothendieck, Produits tensoriels topologiques et espaces nucléaires, *Mem. Am. Math. Soc.* 16 (1955).
- [16] E. Hille, Remarks on ergodic theorems, *Trans. Am. Math. Soc.* 57 (1945) 246–269.
- [17] T. Kalmes, Power bounded weighted composition operators on function spaces defined by local properties, *J. Math. Anal. Appl.* 471 (2019) 211–238.
- [18] T. Kalmes, Dynamics of weighted composition operators on function spaces defined by local properties, *Stud. Math.* 249 (2019) 259–301.
- [19] U. Krengel, *Ergodic Theorems*, De Gruyter Studies in Mathematics, vol. 6, Walter de Gruyter Co., Berlin, 1985.
- [20] A. Przewacki, Dynamical properties of weighted composition operators on the space of smooth functions, *J. Math. Anal. Appl.* 445 (2017) 1097–1113.
- [21] L. Schwartz, *Théorie des distributions, nouvelle édition*, in: Publications de l’Institut de Mathématique de l’Université de Strasbourg, No. IX-X, 1966.
- [22] K. Yosida, *Functional Analysis*, 6th ed., Springer, Berlin, 1980.